


INFECTION CONTROL IN THE HOME OF PATIENTS



After **surgery**, the point of incision is a potential point of entry for germs.



Germs can be transmitted through direct contact from person to person.



Germs can be transmitted through direct contact with a source of infection.



Before starting any **procedure** always **wash** your hands **thoroughly**.



Always wear **disposable gloves** to **avoid** contact with **soiled linens** or if you have a **wound** yourself.



Always **wipe up spills** from any surfaces you have been working on or around.



Discard gloves, **wipes**, **towels** or **pads** you used into a **container**.



In the bathroom, toilets are disinfected with **bleach**.



Liquid **soap** is preferred to bar soap which can be an **asylum** for germs.



Bed linens that are soiled with body fluid are placed into a plastic **bag** to be **carried** to the washing machine.

surgery
procedure
(to) wash
thoroughly
disposable
gloves
(to) avoid
soiled linens
wound
(to) wipe up
spills
(to) discard

opération chirurgicale
acte
laver
minutieusement / à fond
jetable
gants
éviter
linge sale
plaie
essuyer
éclaboussures
jeter

wipes	lingettes
towel	serviette
pad	compresse
container	conteneur
bleach	eau de javel
soap	savon
asylum	nid
bag	sac
(to) carry	porter, transporter